



Generating QCD Gauge Configurations

The Fourth International Workshop on Numerical Analysis and Lattice QCD

Yale University

Michael Clark
Boston University





Talk Outline

- Introduction
- Hybrid Monte Carlo
- Algorithm Improvement
- “Non-local” Algorithms
- Conclusion





Introduction

- Lattice QCD path integral

$$\langle \Omega \rangle = \frac{1}{Z} \int [dU] e^{-S_g(U)} [\det \mathcal{M}(U)]^\alpha \Omega(U)$$

$\alpha = \frac{N_f}{2}$ ($\frac{N_f}{4}$) for Wilson (staggered) fermions, $\mathcal{M} = M^\dagger M$

- $10^8 - 10^9$ degrees of freedom \Rightarrow Monte Carlo integration
- Interpret $e^{-S_g} \det \mathcal{M}^\alpha$ as a Boltzmann weight, and use importance sampling


$$\langle \Omega \rangle \approx \frac{1}{N} \sum_{i=1}^N \Omega(U_i)$$





Quenched Approximation

- Fermion determinant extremely non-local object
- Quenched approximation: set $\det \mathcal{M} = 1$
- Gauge action local: over relaxed heatbath algorithms very efficient
- Just plain wrong!



⇒ Must include dynamical fermions to obtain QCD





The HMC Algorithm (Duane *et al*)

- *De facto* algorithm for including dynamical fermions
- Rewrite determinant in terms of pseudo-fermions

$$\det \mathcal{M} = \int D\phi^\dagger D\phi e^{-\phi^\dagger \mathcal{M}^{-1} \phi} = \int D\phi^\dagger D\phi e^{-S_f}$$

- Need global updates since pseudofermion action is non-local
- Introduce fictitious momentum field π and define a Hamiltonian

$$H = \frac{1}{2} \text{tr} \pi^2 + S_g + S_f = T + S$$

- Integrate Hamilton's equations to propose a new configuration
- Global Accept / Reject to obtain desired probability distribution

$$P(U, \phi) = \frac{1}{Z} e^{-S_g - S_f}$$





The HMC Algorithm (Duane *et al*)

- Each update consists of
 - Hybrid Molecular Dynamics Trajectory
 - * Momentum refreshment heatbath ($P(\pi) \propto e^{-\pi^* \pi / 2}$).
 - * Pseudo-fermion heatbath ($\phi \propto M^\dagger \xi$, where $P(\xi) \propto e^{-\xi^* \xi}$).
 - * MD trajectory with $\tau / \delta\tau$ steps.
 - Metropolis Acceptance Test $P_{\text{acc}} = \min(1, e^{-\delta H})$





Molecular Dynamics

- Hamilton's equations $\frac{dU}{d\tau} = \frac{dT}{d\pi} = \pi$ and $\frac{d\pi}{d\tau} = -\frac{dS}{dU} = F$
- Must discretize the "fictitious time" τ and integrate numerically
- Define integrators in terms of evolution operators Q and P

$$Q \equiv \frac{dT}{d\pi} \frac{\partial}{\partial U} \quad \text{with} \quad e^{\delta\tau Q} : f(U, \pi) \rightarrow f(U + \delta\tau T'(\pi), \pi)$$

$$P \equiv -\frac{dS}{dU} \frac{\partial}{\partial \pi} \quad \text{with} \quad e^{\delta\tau P} : f(U, \pi) \rightarrow f(U, \pi - \delta\tau S'(U))$$

- Metropolis requires Detailed Balance
 - Integration must be reversible and area preserving
 - Use Symmetric Symplectic Integrators, e.g., leapfrog

$$U(\delta\tau)^{\tau/dt} = \left(e^{\delta\tau P/2} e^{\delta\tau Q} e^{\delta\tau P/2} \right)^{\tau/\delta\tau} + O(\delta\tau^2)$$





Molecular Dynamics Forces

- Pure Gauge Force ($\frac{dS_g}{dU}$) local analytic quantity
 - CHEAP
- Fermion force

$$\frac{dS_f}{dU} = \frac{d}{dU} \phi^\dagger \mathcal{M}^{-1} \phi = -\phi^\dagger \mathcal{M}^{-1} \frac{d\mathcal{M}}{dU} \mathcal{M}^{-1} \phi$$


- Each update to the momentum requires solution to $\mathcal{M}\chi = \phi$
- Generally calculated using a Krylov solver, e.g., CG
- EXPENSIVE





Cost of HMC

- Condition number blows up as $m \rightarrow 0$
- Force $\propto 1/m$, requires $\delta\tau \rightarrow 0$ to maintain acceptance rate
- Also, as $m \rightarrow 0$, correlation lengths diverge
- $C \propto \left(\frac{m_\pi}{m_\rho}\right)^{-6} L^5 a^{-7}$
CP-PACS and JLQCD, 2002



⇒ Require huge computers OR better algorithms





Chronological Inverter

- Solution $\chi(\tau) = \mathcal{M}(\tau)^{-1}\phi$ is a smooth function
- Idea: Use previous solutions to act as an initial guess
- Minimize over the space of previous solutions (*Brower et al*) :
for

$$x_0 = \sum_i c_i \mathcal{M} \chi_i$$

solve

$$\chi_j^\dagger \phi = \sum_i c_i \chi_j^\dagger \mathcal{M} \chi_i$$

- Requires high precision solutions to maintain reversibility
- Gain around a factor of 2





Higher Order Integrator

- Potential for gain from using $O(\delta\tau^4)$ integrator, e.g., Campostrini

$$U(\delta\tau)^{\tau/dt} = \left(e^{\delta\tau \epsilon P/2} e^{\delta\tau \epsilon Q} e^{\delta\tau (1-\sigma)P/2} e^{-\delta\tau \epsilon\sigma Q} e^{\delta\tau (1-\sigma)P/2} e^{\delta\tau \epsilon Q} e^{\delta\tau \epsilon P/2} \right)^{\tau/\delta\tau} + O(\delta\tau^4)$$

- Better volume scaling $V^{9/8}$ vs. $V^{5/4}$
- Constructed from sub-leapfrog steps with $\delta\tau^{sub} > \delta\tau$
- Sub-leapfrog integrator can go unstable much sooner than $\delta\tau$ suggests (Joó *et al*)
- Higher order integrators are very unstable
- $O(\delta\tau^2)$ usually found to be optimal





Multiple Timescale Integration (Sexton/Weingarten, 1992)

- For Hamiltonians of the form $H = T + S_1 + S_2$
- Integrate S_1 and S_2 force contributions on different timescales

$$U(\delta\tau)^{\tau/\delta\tau} = \left(\left(e^{\delta\tau P_1/4m} e^{\delta\tau Q/2m} e^{\delta\tau P_1/4m} \right)^m e^{\delta\tau P_2} \left(e^{\delta\tau P_1/4m} e^{\delta\tau Q/2m} e^{\delta\tau P_1/4m} \right)^m \right)^{\tau/\delta\tau}$$

- Two separate timescales $\delta\tau^{S_1} = \delta\tau/m$, $\delta\tau^{S_2} = \delta\tau$
- Large and cheap force = P_1 , Small and expensive force = P_2 ,
- Naïve partitioning: $S_1 = S_g, S_2 = S_f$ fails as $m \rightarrow 0$
- Can extend this recursively for N timescales

$$H = T + S_1 + \dots + S_N$$

- Great idea, but dormant for 10 years...





Multiple Pseudofermions with Mass Preconditioning

- Mass-precondition the fermion determinant (Hasenbusch)

$$\det(M^\dagger M) = \det(\widehat{M}^\dagger \widehat{M}) \det(\widehat{M}(M^\dagger M)^{-1} \widehat{M}^\dagger)$$

with $m(\widehat{M}) > m(M)$



- Tune $\kappa(\widehat{M}^\dagger \widehat{M}) \approx \kappa(\widehat{M}(M^\dagger M)^{-1} \widehat{M}^\dagger) \approx \sqrt{\kappa(M^\dagger M)}$ (Hasenbusch-Jansen)
- Why does this work?
 - Better sampling of Gaussian integral using multiple pseudo-fermions
 - This reduces fluctuations in the fermion force
 - Fermion force $F \propto \kappa^\nu$
- Factor 2 improvement through $\delta\tau$ increase
- Use more than one dummy operator
- Gain increases as $m_l \rightarrow 0$






Multi-timescale Mass Preconditioning (QCDSF, Urbach *et al*)

- Tune dummy operators so that most expensive (preconditioned) force contributes the least
- Use multi-timescale integrator ($S_1 = S_g$, $S_2 = \hat{S}$, $S_3 = S_f$)
- Factor 10 improvement at light quark mass





Domain Decomposition (Lüscher)



- Can rewrite the Dirac operator $M = \begin{pmatrix} D_{\Omega} & D_{\delta\Omega} \\ D_{\delta\Omega'} & D_{\Omega'} \end{pmatrix}$
 - D_{Ω} ($D_{\Omega'}$) is Dirac operator on black (white) blocks with Dirichlet boundaries
 - $D_{\delta\Omega}$ and $D_{\delta\Omega'}$ is the Dirac operator connecting these blocks
 - Rewrite determinant
- $$\det M = \det \begin{pmatrix} D_{\Omega} & 0 \\ 0 & D_{\Omega'} \end{pmatrix} \det \begin{pmatrix} D_{\Omega} & D_{\delta\Omega} \\ D_{\delta\Omega'} & D_{\Omega'} \end{pmatrix} \begin{pmatrix} D_{\Omega}^{-1} & 0 \\ 0 & D_{\Omega'}^{-1} \end{pmatrix}$$
- Separation into large and cheap, and small but expensive
 \Rightarrow multi timescale integrator
 - Large speed up over naive algorithm





Determinant Preconditioning is Key

- Other methods
 - U.V. Filtering (de Forcrand)
 - Polynomial filtering (Peardon and Sexton)
 - Multistep stochastic correction (see talk by Enno)
- All improvements rely on determinant preconditioning





Non-local Actions

- Strange quark inclusion requires $\det \mathcal{M}^{\frac{1}{2}}$
- Finite temperature calculations typically use staggered quarks
 - Remnant chiral symmetry important here
 - Non-local action: $\det \mathcal{M}^{\alpha}$, $\alpha = \frac{1}{2}, \frac{1}{4}$
- HMC cannot be applied for these cases
- Inexact algorithms traditionally used
- Is using exact algorithms more expensive?
- Are the improvements in “Local actions” applicable here?





The R Algorithm (Gottlieb *et al*)

- Rewrite fermionic determinant:

$$\det \mathcal{M}^\alpha = \exp(\alpha \operatorname{tr} \ln \mathcal{M}) = \exp(-S_{\text{eff}})$$

- Integrate Hamilton's equations as before
- Use noisy estimator ξ for trace \equiv pseudo-fermion force
- Leading order error term $O(\delta\tau)$!
- Recover $O(\delta\tau^2)$ with N_f dependent ξ updating
 - Non-reversible
 - Jacobian $\neq 1$
- Cannot include Metropolis acceptance test
 - \Rightarrow Algorithm is inexact
- Naïve cost = HMC, but requires extrapolation to zero stepsize
- Stepsize rule of thumb $\delta\tau \sim \frac{2}{3}m_1$





Polynomial Hybrid Monte Carlo (de Forcrand-Takaishi, Frezzotti-Jansen)

- Write in pseudo-fermion notation

$$\begin{aligned} \det \mathcal{M}^\alpha &= \int \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{-\bar{\psi} \mathcal{M}^{-\alpha} \psi} \\ &\approx \int \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{-\bar{\psi} P(\mathcal{M}) \psi}, \end{aligned}$$


where $P(\mathcal{M})$ is valid over spectrum

- Pseudo-fermion heatbath easily realised $P(\mathcal{M}) = p^\dagger(\mathcal{M})p(\mathcal{M})$
- Use standard MD leapfrog \Rightarrow exact
- Generally polynomial degree $m > N_{\text{iter}}$ CG iterations
- Use low degree polynomial
 - Reweight acceptance test or observable
- Use high degree polynomial
 - $\alpha \neq 1$ derivative uses Leibniz rule \Rightarrow Memory, rounding





Optimal rational approximations



- Generated using Remez algorithm
- Real non-degenerate roots (poles are always +ve)
- Partial fractions - $r(x) = \sum_{k=1}^m \frac{\alpha_k}{x + \beta_k}$
- Evaluate using multi-shift solver
- Numerically stable (α_k have same sign)





Rational Hybrid Monte Carlo (Clark-Kennedy)

- Rewrite fermionic determinant

$$\begin{aligned} \det \mathcal{M}^\alpha &= \int \mathcal{D}\bar{\phi} \mathcal{D}\phi e^{-\bar{\phi} \mathcal{M}^{-\alpha} \phi} \\ &\approx \int \mathcal{D}\bar{\phi} \mathcal{D}\phi e^{-\bar{\phi} r^2(\mathcal{M}) \phi}, \end{aligned}$$

with $r(x) = x^{-\alpha/2}$

- Precision is cheap: Conventional Metropolis
- RHMC:
 - Hybrid Molecular Dynamics Trajectory
 - * Momentum refreshment heatbath ($P(\pi) \propto e^{-\pi^* \pi/2}$).
 - * Pseudo-fermion heatbath ($\phi \propto r(\mathcal{M})^{-1} \xi$, where $P(\xi) \propto e^{-\xi^* \xi}$).
 - * MD trajectory with $\tau/\delta\tau$ steps.
 - Metropolis Acceptance Test $P_{\text{acc}} = \min(1, e^{-\delta H})$





Rational Hybrid Monte Carlo (Clark-Kennedy)

- MD trajectory
 - Double inversion from $r^2(\mathcal{M})$
 - Use low degree approx $\bar{r} \approx \mathcal{M}^{-\alpha} \approx r^2$
 - Pseudo-fermion force

$$S'_{\text{pf}} = - \sum_{i=1}^{\bar{m}} \bar{\alpha}_i \phi^\dagger (\mathcal{M} + \bar{\beta}_i)^{-1} \mathcal{M}' (\mathcal{M} + \bar{\beta}_i)^{-1} \phi.$$


- CG cost per trajectory \approx HMC
 - One extra inversion required for heatbath






Exact vs. Inexact (RBC-Bielefeld)

RHMC vs. HMDR, p4fat3 $m_q=0.01, 8^3 \times 4$



RHMC vs. HMDR, p4fat7, $m_q=0.1, 8^3 \times 4$





- P4 staggered fermions
- RHMC allows an $O(10)$ increase in stepsize
- Speedup greater as $m_l \rightarrow 0$





Exact vs. Inexact (de Forcrand-Philipsen)



(Naïve Staggered Fermions, $N_f = 3$, $V = 8^3 4$)

- Results:
 - Binder cumulant increases
 - Stepsize extrapolation is vital for R algorithm
 - 25% reduction in critical quark mass at $\delta\tau^R = \frac{1}{2}m_l$
 - 20% change in renormalised quark mass
- Conclusion: “an exact algorithm is mandatory”





Multiple Pseudofermions with RHMC (Clark-Kennedy)

- Rewrite determinant


$$\begin{aligned} \det \mathcal{M} &= [\det \mathcal{M}^{1/n}]^n \\ &\propto \prod_{j=1}^n d\phi_j d\phi_j^\dagger \exp\left(-\phi_j^\dagger \mathcal{M}^{-1/n} \phi_j\right), \end{aligned}$$

- So called n^{th} root trick
- Speedup through $\delta\tau$ increase
- No dummy mass parameters to tune \Rightarrow easy to increase n
- Single fermion timescale





Integrator Instability (Clark-Kennedy)




(Staggered fermions, $V = 16^4$, $\beta = 5.6$, $N_f = 2$, $m = 0.005$)

- With $n = 1$ integrator breaks down as $\delta\tau$ is increased
- Instability “tickled” by low fermion modes $\sim O(\frac{1}{m_l})$ (Joó *et al*)
- Higher order integrators are more ticklish
- What happens with multiple pseudofermions?





Integrator Instability (Clark-Kennedy)



(Staggered fermions, $V = 16^4$, $\beta = 5.6$, $N_f = 2$, $m = 0.005$)

- Removes instability in the integrator!
- Why does this work? Lowest modes now $O(\frac{1}{m_1})^{1/n}$
- Force now bulk dominated
- Higher order integrators now beneficial





Who's the fastest of them all? ($N_f = 2$ Wilson)

- Compare multi-timescale mass preconditioning and high order RHMC
- Use popular testing parameters ($V = 24^3.32$, $\beta = 5.6$, Plaquette + Wilson fermions)
- Use measure $C = \tau_{\text{int}}^{\text{plaq}} \cdot N_{\text{mv}} \cdot 10^{-4}$

	C		
κ	RHMC	Urbach <i>et al</i>	Orth <i>et al</i>
0.15750	9.6	9.0	19.1
0.15800	29.9*	17.4	128
0.15825	52.5*	56.5	-

*Using 4MN5 fourth order integrator (de Forcrand and Takaishi)

- RHMC similar in cost to mass preconditioning
- Look at integrator stability with mass preconditioning
 - Gain from higher order integrators also?





2+1 QCD

- e.g. Domain Wall 2+1 flavour determinant

$$\left(\frac{\det M_l^\dagger M_l}{\det M_{pv}^\dagger M_{pv}} \right) \left(\frac{\det M_s^\dagger M_s}{\det M_{pv}^\dagger M_{pv}} \right)^{1/2} = \left(\frac{\det M_l^\dagger M_l}{\det M_s^\dagger M_s} \right) \left(\frac{\det M_s^\dagger M_s}{\det M_{pv}^\dagger M_{pv}} \right)^{3/2}$$

- Mass Precondition using the strange quark
- Use n^{th} root trick for triple strange
- NOT mutually exclusive improvements
- Use multi-timescale integrator (gauge, triple strange, light)
- Light quark mass constitutes around 10% CG cost
- Cost dependence on mass comes mostly from autocorrelation








Berlin Wall Plot

- Compare the cost of fermion formulations and/or algorithms
 - $N_f = 2 + 1$ DWF RHMC (RBC-UKQCD)
 - $N_f = 2$ Mass preconditioned Wilson (Urbach *et al*)
 - $N_f = 2$ Mass preconditioned Clover (QCDSF)
 - $N_f = 2 + 1$ Mass preconditioned Clover + RHMC (Wuppertal-Jülich)
 - $N_f = 2$ Mass preconditioned Twisted Mass (ETM)
 - $N_f = 2 + 1$ AsqTad R (MILC)
 - $N_f = 2 + 1$ AsqTad RHMC (Clark-Kennedy)
- All data scaled to $V = 24^3 \times 40$, $a = 0.08$
- Box is too small and too coarse











Conclusions

- After 20 years HMC is still the best dynamical algorithm
- The last 5 years has seen an explosion in HMC improvement
- Determinant preconditioning is the key behind all improvement
- Non-local actions are no problem
- Pick and mix the most appropriate algorithm
 - Multiple time scale mass preconditioning
 - Domain decomposition
 - n^{th} rootary with RHMC
- HMC cost is now $C \propto \left(\frac{m_\pi}{m_\rho}\right)^{-2} L^5 a^{-6}$ (Lüscher)
- Physical point no longer a pipedream
- Further improvement must come from autocorrelation?

